



ELSEVIER

ScienceDirect

Current Opinion in
Plant Biology

Plant carotenoids: genomics meets multi-gene engineering

Giovanni Giuliano



Carotenoids are present in plant photosynthetic tissues, where they have essential roles in photoreception and photoprotection, as well as in non-photosynthetic tissues, where they act as colorants, precursors for plant isoprenoid volatiles and signaling molecules (abscisic acid and strigolactones), nutritional antioxidants and vitamin A precursors. This review presents the recent advances in our understanding of their biosynthesis, the key metabolic steps controlling their accumulation in plant non-photosynthetic tissues and their metabolic engineering using multi-gene approaches.

Addresses

Italian National Agency for New Technologies, Energy and Sustainable Development (ENEA), Casaccia Research Center, Via Anguillarese 301, Roma 00123, Italy

Corresponding author: Giuliano, Giovanni (giovanni.giuliano@enea.it)

Current Opinion in Plant Biology 2014, **19**:111–117

This review comes from a themed issue on **Physiology and metabolism**

Edited by **Sarah E O'Connor** and **Thomas P Brutnell**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 7th June 2014

<http://dx.doi.org/10.1016/j.pbi.2014.05.006>

1369-5266/© 2014 The Author. Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

Introduction

Carotenoids are C40 isoprenoid compounds, made of C5 isopentenyl building blocks, which in plants are synthesized through two separate pathways: the cytoplasmic mevalonate pathway and the plastid-located (or MEP) pathway (Figure 1a). The MEP pathway provides a large part of the carbon flux that goes into carotenoids [1]. Its main products, isopentenyl pyrophosphate (IPP) and dimethylallyl pyrophosphate (DMAPP), are condensed into geranylgeranyl pyrophosphate (GGPP), which acts as a precursor for a series of isoprenoid molecules, including carotenoids (Figure 1a). The condensation of two GGPP molecules yields 15-*cis*-phytoene, a colorless, linear carotene, which is converted into lycopene by a poly-*cis* desaturation/isomerization pathway relying on two desaturases and two isomerases. Lycopene is the substrate of two competing cyclases, introducing β -rings and ϵ -rings at the ends of the molecule. A series of heme and non-heme hydroxylases, one epoxidase and one de-epoxidase,

complete the canonical plant pathway. Some steps (the biosynthesis of neoxanthin, the 9-*cis* isomerization of neoxanthin and violaxanthin) await complete elucidation. An excellent review on the mechanistic aspects of carotenoid biosynthesis was published recently [2].

Carotenoids are the substrates of cleavage enzymes, which come in at least two types (Figure 1): nine-*cis*-epoxycarotenoid dioxygenases (NCEDs) that catalyze the first step in abscisic acid (ABA) biosynthesis, and carotenoid cleavage dioxygenases (CCDs) that catalyze a vast array of different cleavage steps giving rise to both volatile and non-volatile apocarotenoids, with functions as aromas, allelopathic substances or hormone-like substances (strigolactones) [3].


Metabolic engineering of plant carotenoid biosynthesis has focused primarily on two targets. One target is β -carotene, a nutritional vitamin A precursor, for alleviating vitamin A deficiency in populations whose diet is deficient in this vitamin. These biofortification efforts, utilizing both transgenic and classical breeding approaches, have focused on staple foods that are deficient in β -carotene, such as rice, white maize, potato and cassava [4[•],5[•],6,7^{••},8[•],9[•],10[•]].

Additional targets are high value xanthophylls used as food and feed additives. Certainly the most explored one is astaxanthin, a dihydroxy, diketo xanthophyll (Figure 1) synthesized by many marine bacteria and algae and by a handful of plants and fungi, used both as a dietary antioxidant and as a feed additive in fisheries. Astaxanthin is presently produced through chemical synthesis or extraction from microalgae, and alternative sources are needed to meet market demand at competitive prices. Transgenic and transplastomic approaches for the production of astaxanthin *in planta* have focused on tobacco, carrot, tomato, potato and maize [7^{••},11[•],12,13,14,15[•],16[•],17,18].

Updating the carotenoid pathway

In organisms with oxygenic photosynthesis (cyanobacteria and plants) the conversion of 15-*cis*-phytoene to lycopene is effected through a poly-*cis* pathway, involving the action of two desaturases introducing *cis* double-bonds and two isomerases converting them back to the all-*trans* configuration (Figure 1a). Phytoene desaturase (PDS), ζ -carotene desaturase (ZDS) and prolycopene isomerase (CrtISO) are flavoproteins catalyzing non-redox reactions [19,20], like the bacterial CrtI enzyme, which is however able to perform all the reactions

Figure 1



required for lycopene synthesis [21[•]]. This catalytic versatility of *CrtI* has prompted its use in many metabolic engineering efforts. A second plant isomerase, ζ -carotene isomerase or *ZISO*, able to isomerize the 15-*cis* double bond, was recently isolated [22^{••}]. *ZISO* encodes a small protein, very different from *CrtISO*, and contains consensus sequences for iron binding.

The biosynthesis of astaxanthin from β -carotene in marine bacteria was elucidated over 15 years ago [23^{••}]. It is mediated by two genes, *CrtZ* and *CrtW*, encoding a hydroxylase and a ketolase, respectively, acting in a combinatorial fashion on the two β -rings (Figure 1b). The pathway for astaxanthin biosynthesis in flowers of the higher plant *Adonis aestivalis* was recently elucidated [24^{••}] (Figure 1c): first, a β -ring 4-dehydrogenase (CBFD) catalyzes the formation of a 4-hydroxyl group, which is then converted to a 4-keto group by a hydroxy- β -ring 4-dehydrogenase (HBFD); finally, CBFD inserts a 3-hydroxyl group, completing the biosynthesis. This pathway has been shown to be active in *Escherichia coli*; in contrast to the bacterial enzymes, which seem to act in a competitive fashion when modifying the same ring, leading to the accumulation of intermediates [25], the plant enzymes seem to act synergistically, leading to complete conversion of β -carotene into astaxanthin. Despite this, to date there are no examples of transfer of the *Adonis* pathway to other flowering plants.

Genomics comes of age

Potato and tomato have been extensively utilized in the past for metabolic engineering efforts. Sequencing of their genomes [26[•],27^{••}] unveiled several novel genes in carotenoid biosynthesis and their regulation. In both plants, the first dedicated enzyme of the pathway, phytoene synthase (PSY) (Figure 1a), is encoded by three genes, two of which (*PSY1* and *PSY2*) show expression in most tissues, while expression of *PSY3* is very low and confined to roots. The *PSY1*–*PSY2* pair was generated by the *Solanum* whole genome triplication dating back to approximately 70 million years ago (mya), while the origin of *PSY3* seems more ancient [27^{••}]. The expression pattern of the *Solanum PSY1* remained generally similar to that of its *PSY2* counterpart until recently (0.5–1 mya), when its overexpression in fruits gave rise to the lycopene accumulation pattern characteristic of the red fruits of tomato and *S. pimpinellifolium* (G Falcone and G Giuliano, unpublished data). *PSY* and *ZISO* are the two most highly expressed genes in the carotenoid pathway in fruits of red-fruited species, while they are expressed at low levels in fruits of green-fruited species (G Falcone and G Giuliano, unpublished data). *PSY1* is absolutely necessary for lycopene accumulation in tomato fruits, while the

function of *ZISO* can be partially substituted by light [28[•]].

PSY is a key regulator of carotenoid accumulation in staple crops such as maize [29[•]], durum wheat [30] and cassava [10[•]]. A single nucleotide polymorphism in a *PSY* gene highly expressed in cassava roots co-segregates with high β -carotene levels. This polymorphism results in a single amino acid change in a highly conserved region of the protein which, when tested in *E. coli*, results in increased catalytic activity. A systematic comparison showed that *PSY* genes from different plant sources differed greatly in their capacity to induce accumulation of β -carotene in the endosperm of golden rice, with rice and maize *PSY* being the most efficient [5[•]]. A recent survey of several *PSY* enzymes from different sources showed that the majority of them localize to plastoglobules, with the exception of maize *PSY1*, which localizes preferentially in the stroma [31[•]].

PSY seems to be a key integrator for several signals regulating the carotenoid pathway. In *Arabidopsis*, the phytochrome-interacting transcription factor PIF1 directly interacts with the promoter of the *PSY* gene, mediating its light-regulated transcription [32[•]]. In tomato, the MADS box regulator of ripening, RIN, interacts with the *PSY* promoter [33,34,35]. Overexpression in potato tubers of the cauliflower Or regulator, a molecular chaperone able to induce formation of chromoplasts in heterologous systems [36], results in increased *PSY* stability [37]. The tomato Stay Green SISGR1 protein regulates fruit lycopene and β -carotene accumulation through direct interaction with the *PSY* enzyme [38]. Finally, it has been proposed that epistasis in tomato color mutations involves regulation of *PSY1* expression by *cis*-carotenoids [39[•]].

β -carotene hydroxylase (CHY) and lycopene ϵ -cyclase (LCY-*e*) are important regulators of β -carotene and total carotenoid accumulation in maize. Mutations at the *LCY-e* locus redirect the flux towards the β -branch, increasing β -carotene content [40[•]]; *CHY* alleles associated with reduced transcript expression correlate with higher β -carotene concentrations [9[•],41]. In potato, transgenic silencing of the *CHY* and *LCY-e* genes results in increased β -carotene and total carotenoid content [42,43]. A dominant allele of *CHY2* is associated with accumulation of carotenoids, especially β -xanthophylls, in potato [44[•]] and the *CHY2* gene is more expressed in the high-carotenoid DM potato genotype than in the low carotenoid RH one [26[•]]. A second gene mainly controlling the accumulation of zeaxanthin in potato is zeaxanthin epoxidase (*ZEP*) (Figure 1). Loss-of-function alleles of *ZEP*,

(Fig.1 legend continue) (a) Schematic carotenoid biosynthetic pathway. (b) Astaxanthin biosynthesis in marine bacteria [23^{••}]. (c) Astaxanthin biosynthesis in *Adonis aestivalis* [24^{••}]. Names of compounds are in black boldface. Names of enzymes are black (plant enzymes) and red (bacterial enzymes).

or silencing of the *ZEP* gene, cause an accumulation of zeaxanthin [44•,45].

Sequencing of the peach genome [46•] helped clarify the molecular basis of the high-carotenoid, yellow flesh trait in peach. White flesh is dominant over yellow in peach, and is associated with increased emission of isoprenoid volatiles, likely derived from the cleavage of carotenoid molecules, suggesting that the trait is due to a gain-of-function mutation in a carotenoid cleavage dioxygenase. Analysis of a yellow-fleshed cultivar and of its white-fleshed mutant showed that a *CCD4* transcript was induced in the mutant [47•]. Analysis of a larger genotype collection suggested that the white flesh color (active *CCD4*) was the ancestral situation in peach and that yellow-fleshed genotypes arose in modern cultivars through three independent mutational mechanisms (nucleotide substitutions, small insertions and transposable element insertions) [48•].

CCD4 is also a key determinant of carotenoid pigmentation in petals of *Chrysanthemum* [49] and of *Lilium* [50]. In contrast, in *Calendula* petals *CrtISO* is a key determinant [51•]. *CCD4* is also a key determinant of carotenoid/apocarotenoid content in potato tubers [52], *Arabidopsis* seeds and senescing leaves [53•] and *Citrus* fruits [54•]. In *Arabidopsis* and potato, *CCD4* levels correlate inversely with pigmentation, while in *Citrus* the opposite is true. This is because *Citrus* *CCD4* asymmetrically cleaves β -carotene and zeaxanthin at the 7,8 position, giving rise to non-volatile C30 apocarotenoids (Figure 1a), which accumulate in the flavedo of fruits and contribute to their pigmentation. In potato, RNAi inhibition of *CCD4* caused not only accumulation of carotenoids, but also a series of developmental phenotypes [52], suggesting the intriguing hypothesis that a *CCD4* cleavage product may have a signaling role, or that *CCD4* silencing in tubers re-routes the flux towards other cleavage pathways with a signaling role (NCED, giving rise to ABA and/or CCD7/8, giving rise to strigolactones).

Multiple gene engineering

The stable overexpression of multiple transgenes is one of the main bottlenecks limiting reconstruction of complex biochemical pathways. Early examples of mini-pathway engineering in rice, tomato and potato employed 2–3 transgenes cloned in a single *Agrobacterium* construct [4•,5•,6,55]. More recently, up to seven different bacterial carotenoid genes (from isopentenyl pyrophosphate isomerase (*IP1*) to *CrtW*, Figure 1), have been assembled in a single *Agrobacterium* construct and used to transform canola and *Lilium* [56,57].

A second approach for mini-pathway engineering is combinatorial transformation of independent gene constructs. Five independent transgenes, from *PSY* to *CrtW* (Figure 1), were introduced in a white endosperm maize

using biolistic transformation [7••]. Different combinations of expressed transgenes, from none to all five, were recovered in an approximately normal distribution. This approach was brought one step further by stacking two carotenoid genes and one gene each from the ascorbate and folate pathways to obtain maize simultaneously fortified with β -carotene (provitamin A), vitamin C and folate [8•].

A third approach is the expression of different genes (*CrtB* and *CrtI*) as a single transcript under the control of a single promoter [58•]. The downstream gene product is expressed either by inserting an internal ribosomal entry site (IRES) from a plant virus, or by inserting between the two proteins a self-cleaving protein sequence (2A) from an animal virus. The polyprotein construct is ninefold more efficient than the IRES construct in rice endosperm [58•] and has been used to increase total carotenoid and β -carotene levels in soybean seeds [59].

A fourth approach, applicable in principle to any plastid-localized biosynthetic pathway, is the expression of multiple genes as a single operon inserted in plastid DNA. This approach can be used to insert biosynthetic pathways from a different organism, or from a different cell compartment, to plastids: the *CrtZ* and *CrtW* genes for astaxanthin biosynthesis have been overexpressed in tobacco [15•] and lettuce [17]. In a separate approach, the whole cytoplasmic mevalonate pathway for biosynthesis of IPP was transferred to tobacco plastids [60•]. The functionality of the transferred pathway was demonstrated through resistance to fosmidomycin (an inhibitor of the MEP pathway). Transplastomic plants accumulated higher levels of carotenoids, sterols and triacylglycerols than control plants.

In general, the carotenoids accumulated in plants engineered with multiple genes are consistent with the function of the introduced transgenes, with a series of interesting exceptions:

- (i) The multifunctional *CrtI* desaturase/isomerase utilized in many metabolic engineering experiments is unable to catalyze complete conversion of phytoene into downstream compounds [6,7••], suggesting that part of the phytoene pool is inaccessible to *CrtI*. Fractionation of transgenic tomato chromoplasts overexpressing *CrtB* + *CrtI* showed that phytoene was mainly found in plastoglobules, whereas the *CrtB* and *CrtI* enzymes were localized to the membrane fraction [61•]. A series of enzymes for carotenoid metabolism are known to localize to plastoglobules, including ZDS, LCY-b, CHY and *CCD4* [62,63], whereas *PSY1* has been reported to be localized in both plastoglobules and stroma [31•,61•].

- (ii) A lycopene β -cyclase (*LCY-b*) transgene is not required for β -carotene accumulation in different plant tissues (tomato fruits and maize and rice endosperm). All three tissues overexpressing *CrtB/PSY + CrtI* accumulate significant levels of β -carotene, [4^{••}, 7^{••}, 61[•]]. Endogenous *LCY-b* expression is probably the cause of this phenotype.
- (iii) In plants transformed with multiple bacterial genes for the production of astaxanthin, the biosynthesis is often arrested at intermediate compounds, such as echinenone, 3'-OH-echinenone, canthaxanthin and adonixanthin [7^{••}, 56, 57] (Figure 1b). This limitation can be overcome by the use of algal ketolases and hydroxylases showing co-operative, rather than antagonistic action [18, 64, 65] or by plastid transformation [15[•], 17].

Concluding remarks

In recent years, the availability of genomic tools, including whole genome sequences, has uncovered a series of quantitative trait loci (QTLs) controlling carotenoid accumulation in agricultural plants. These QTLs can now be used in breeding programs aimed at optimizing plant carotenoid composition. At the same time, advanced tools for engineering of the nuclear and plastid genomes have been developed, including combinatorial nuclear transformation [7^{••}] and artificial *cis*-elements for the expression in chromoplasts [66[•]]. The additional availability of synthetic biology tools [67] and metabolic models accounting for subcellular compartmentalization and tissue-specificity [68] will dramatically improve our capacity to manipulate plant carotenoid biosynthesis for biofortification and green chemistry applications.

Note added in proof

A gene essential for neoxanthin biosynthesis was recently cloned from tomato [69].

Acknowledgements

Work in GG's laboratory is supported by the Italian Ministry of Agriculture (NUTRISOL project) the Italian Ministry of Research ('Integrated Knowledge for the Sustainability and Innovation of Italian Agri-Food' project) and by the European Commission (DISCO project).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Rodriguez-Concepcion M: **Supply of precursors for carotenoid biosynthesis in plants.** *Arch Biochem Biophys* 2010, **504**:118-122.
2. Moise AR, Al-Babili S, Wurtzel ET: **Mechanistic aspects of carotenoid biosynthesis.** *Chem Rev* 2013, **114**:164-193.
3. Auldridge ME, McCarty DR, Klee HJ: **Plant carotenoid cleavage oxygenases and their apocarotenoid products.** *Curr Opin Plant Biol* 2006, **9**:315-321.
4. Ye X, Al-Babili S, Kloti A, Zhang J, Lucca P, Beyer P, Potrykus I: **Engineering the provitamin A (beta-carotene) biosynthetic**

pathway into (carotenoid-free) rice endosperm. *Science* 2000, **287**:303-305.

The original article describing the generation of β -carotene fortified 'Golden rice'.

5. Paine JA, Shipton CA, Chaggar S, Howells RM, Kennedy MJ, Vernon G, Wright SY, Hinchliffe E, Adams JL, Silverstone AL, Drake R: **Improving the nutritional value of Golden Rice through increased pro-vitamin A content.** *Nat Biotechnol* 2005, **23**:482-487.

The generation of 'Golden rice 2' with improved β -carotene content through the use of a highly efficient phytoene synthase.

6. Diretto G, Al-Babili S, Tavazza R, Papacchioli V, Beyer P, Giuliano G: **Metabolic engineering of potato carotenoid content through tuber-specific overexpression of a bacterial mini-pathway.** *PLoS ONE* 2007, **2**:e350.

7. Zhu C, Naqvi S, Breitenbach J, Sandmann G, Christou P, Capell T: **Combinatorial genetic transformation generates a library of metabolic phenotypes for the carotenoid pathway in maize.** *Proc Natl Acad Sci U S A* 2008, **105**:18232-18237.

Combinatorial biolistic transformation of maize with five transgenes: *Z. mays* phytoene synthase 1, *P. ananatis* phytoene desaturase, *G. lutea* lycopene β -cyclase and β -carotene hydroxylase and *Paracoccus* β -carotene ketolase, and the recovery of seven different combinations of transgenes, accumulating different carotenoids. A demonstration of the power of combinatorial transformation.

8. Naqvi S, Zhu C, Farre G, Ramessar K, Bassie L, Breitenbach J, Perez Conesa D, Ros G, Sandmann G, Capell T, Christou P: **Transgenic multivitamin corn through biofortification of endosperm with three vitamins representing three distinct metabolic pathways.** *Proc Natl Acad Sci U S A* 2009, **106**:7762-7767.

An application of the combinatorial transformation concept to the simultaneous biofortification of maize with β -carotene (169-fold), vitamin C (sixfold) and folate (twofold).

9. Yan J, Kandianis CB, Harjes CE, Bai L, Kim EH, Yang X, Skinner DJ, Fu Z, Mitchell S, Li Q et al.: **Rare genetic variation at *Zea mays* *Crtb1* increases beta-carotene in maize grain.** *Nat Genet* 2010, **42**:322-327.

A QTL in the maize *CHY* gene associated with β -carotene concentration in kernels.

10. Welsch R, Arango J, Bar C, Salazar B, Al-Babili S, Beltran J, Chavarriaga P, Ceballos H, Tohme J, Beyer P: **Provitamin A: accumulation in cassava (*Manihot esculenta*) roots driven by a single nucleotide polymorphism in a phytoene synthase gene.** *Plant Cell* 2010, **22**:3348-3356.

A SNP in the cassava *PSY1* gene associated high carotenoids and higher catalytic activity.

11. Mann V, Harker M, Pecker I, Hirschberg J: **Metabolic engineering of astaxanthin production in tobacco flowers.** *Nat Biotechnol* 2000, **18**:888-892.

The first report of metabolic engineering of astaxanthin production in a higher plant.

12. Gerjets T, Sandmann G: **Ketocarotenoid formation in transgenic potato.** *J Exp Bot* 2006, **57**:3639-3645.

13. Morris WL, Ducreux LJ, Fraser PD, Millam S, Taylor MA: **Engineering ketocarotenoid biosynthesis in potato tubers.** *Metab Eng* 2006, **8**:253-263.

14. Zhu C, Gerjets T, Sandmann G: **Nicotiana glauca engineered for the production of ketocarotenoids in flowers and leaves by expressing the cyanobacterial *crtO* ketolase gene.** *Transgenic Res* 2007, **16**:813-821.

15. Hasunuma T, Miyazawa S, Yoshimura S, Shinzaki Y, Tomizawa K, Shindo K, Choi SK, Misawa N, Miyake C: **Biosynthesis of astaxanthin in tobacco leaves by transplastomic engineering.** *Plant J* 2008, **55**:857-868.

The first transplastomic multi-gene engineering of astaxanthin and a demonstration of the capacity of leaves to tolerate the accumulation of unusual carotenoids.

16. Jayaraj J, Devlin R, Punja Z: **Metabolic engineering of novel ketocarotenoid production in carrot plants.** *Transgenic Res* 2008, **17**:489-501.

Carrot roots overexpressing a microalgal ketolase accumulate spectral amounts of astaxanthin and other ketocarotenoids.

17. Harada H, Maoka T, Osawa A, Hattori JI, Kanamoto H, Shindo K, Otomatsu T, Misawa N: **Construction of transplastomic lettuce (*Lactuca sativa*) dominantly producing astaxanthin fatty acid esters and detailed chemical analysis of generated carotenoids.** *Transgenic Res* 2014, **23**:303-315.
18. Huang JC, Zhong YJ, Liu J, Sandmann G, Chen F: **Metabolic engineering of tomato for high-yield production of astaxanthin.** *Metab Eng* 2013, **17**:59-67.
19. Giuliano G, Giliberto L, Rosati C: **Carotenoid isomerase: a tale of light and isomers.** *Trends Plant Sci* 2002, **7**:427-429.
20. Yu Q, Ghisla S, Hirschberg J, Mann V, Beyer P: **Plant carotene cis-trans isomerase CrtISO: a new member of the fad(red)-dependent flavoproteins catalyzing non-redox reactions.** *J Biol Chem* 2011, **286**:8666-8676.
21. Schaub P, Yu Q, Gemmecker S, Poussin-Courmontagne P, Mailliot J, McEwen AG, Ghisla S, Al-Babili S, Cavarelli J, Beyer P: **On the structure and function of the phytoene desaturase crtI from *Pantoea ananatis*, a membrane-peripheral and fad-dependent oxidase/isomerase.** *PLoS One* 2012, **7**:e39550.

A careful mechanistic study on a key enzyme for carotenoid metabolic engineering.

22. Chen Y, Li F, Wurtzel ET: **Isolation and characterization of the ZISO gene encoding a missing component of carotenoid biosynthesis in plants.** *Plant Physiol* 2010, **153**:66-79.

Isolation of a missing isomerase in the plant poly-cis pathway.

23. Misawa N, Satomi Y, Kondo K, Yokoyama A, Kajiwaru S, Saito T, Ohtani T, Miki W: **Structure and functional analysis of a marine bacterial carotenoid biosynthesis gene cluster and astaxanthin biosynthetic pathway proposed at the gene level.** *J Bacteriol* 1995, **177**:6575-6584.

The isolation of a bacterial operon for astaxanthin biosynthesis, expression of the individual genes in *E. coli* and characterization of the synthesized carotenoids by HPLC, NMR and CD spectrometry. A vintage classic.

24. Cunningham FX Jr, Gantt E: **Elucidation of the pathway to astaxanthin in the flowers of *Adonis aestivalis*.** *Plant Cell* 2011, **23**:3055-3069.

Isolation of the missing step in plant astaxanthin biosynthesis through color complementation in *E. coli*. The careful reconstruction of an elusive biochemical pathway.

25. Fraser PD, Shimada H, Misawa N: **Enzymic confirmation of reactions involved in routes to astaxanthin formation, elucidated using a direct substrate in vitro assay.** *Eur J Biochem* 1998, **252**:229-236.
26. Xu X, Pan S, Cheng S, Zhang B, Mu D, Ni P, Zhang G, Yang S, Li R, Wang J *et al.*: **Genome sequence and analysis of the tuber crop potato.** *Nature* 2011, **475**:189-195.

The potato genome sequence. The supplementary materials highlight the genomic basis of high-carotenoid versus low-carotenoid content in tubers.

27. Sato S, Tabata S, Hirakawa H, Asamizu E, Shirasawa K, Isobe S, Kaneko T, Nakamura Y, Shibata D, Aoki K *et al.*: **The tomato genome sequence provides insights into fleshy fruit evolution.** *Nature* 2012, **485**:635-641.

The tomato genome sequence, highlighting how a Solanaceae-specific whole genome triplication generated novel genes controlling fruit characteristics, including PSY1

28. Fantini E, Falcone G, Frusciante S, Giliberto L, Giuliano G: **Dissection of tomato lycopene biosynthesis through virus-induced gene silencing.** *Plant Physiol* 2013, **163**:986-998.

A systematic functional study of nine candidate genes for fruit lycopene biosynthesis using Virus Induced Gene Silencing.

29. Palaisa K, Morgante M, Tingey S, Rafalski A: **Long-range patterns of diversity and linkage disequilibrium surrounding the maize y1 gene are indicative of an asymmetric selective sweep.** *Proc Natl Acad Sci U S A* 2004, **101**:9885-9890.

A selective sweep at the maize Y1 gene, encoding phytoene synthase and controlling kernel pigmentation.

30. Pozniak CJ, Knox RE, Clarke FR, Clarke JM: **Identification of QTL and association of a phytoene synthase gene with endosperm colour in durum wheat.** *Theor Appl Genet* 2007, **114**:525-537.
31. Shumskaya M, Bradbury LM, Monaco RR, Wurtzel ET: **Plastid localization of the key carotenoid enzyme phytoene synthase is altered by isozyme, allelic variation, and activity.** *Plant Cell* 2012, **24**:3725-3741.

An intriguing report on the suborganellar localization of different PSY isozymes.

32. Toledo-Ortiz G, Huq E, Rodriguez-Concepcion M: **Direct regulation of phytoene synthase gene expression and carotenoid biosynthesis by phytochrome-interacting factors.** *Proc Natl Acad Sci U S A* 2010, **107**:11626-11631.

The first molecular link between the phytochrome signal transduction pathway and expression of a key carotenoid gene.

33. Martel C, Vrebalov J, Tafelmeyer P, Giovannoni JJ: **The tomato MADS-box transcription factor ripening inhibitor interacts with promoters involved in numerous ripening processes in a colorless nonripening-dependent manner.** *Plant Physiol* 2011, **157**:1568-1579.
34. Fujisawa M, Nakano T, Ito Y: **Identification of potential target genes for the tomato fruit-ripening regulator rin by chromatin immunoprecipitation.** *BMC Plant Biol* 2011, **11**:26.
35. Fujisawa M, Nakano T, Shima Y, Ito Y: **A large-scale identification of direct targets of the tomato MADS box transcription factor ripening inhibitor reveals the regulation of fruit ripening.** *Plant Cell* 2013, **25**:371-386.
36. Giuliano G, Diretto G: **Of chromoplasts and chaperones.** *Trends Plant Sci* 2007, **12**:529-531.
37. Li L, Yang Y, Xu Q, Owsiany K, Welsch R, Chitchumroonchokchai C, Lu S, Van Eck J, Deng XX, Failla M, Thannhauser TW: **The OR gene enhances carotenoid accumulation and stability during post-harvest storage of potato tubers.** *Mol Plant* 2012, **5**:339-352.
38. Luo Z, Zhang J, Li J, Yang C, Wang T, Ouyang B, Li H, Giovannoni J, Ye Z: **A stay-green protein SISGR1 regulates lycopene and beta-carotene accumulation by interacting directly with SPSY1 during ripening processes in tomato.** *New Phytol* 2013, **198**:442-452.
39. Kachanovsky DE, Filler S, Isaacson T, Hirschberg J: **Epistasis in tomato color mutations involves regulation of phytoene synthase 1 expression by cis-carotenoids.** *Proc Natl Acad Sci U S A* 2012, **109**:19021-19026.

An elegant genetic study explaining a paradox: how a gene located downstream in a biochemical pathway can be epistatic over a gene located upstream.

40. Harjes CE, Rocheford TR, Bai L, Brutnell TP, Kandianis CB, Sowinski SG, Stapleton AE, Vallabhaneni R, Williams M, Wurtzel ET *et al.*: **Natural genetic variation in lycopene epsilon cyclase tapped for maize biofortification.** *Science* 2008, **319**:330-333.

A QTL in the maize LCY-e gene associated with β -carotene concentration in kernels.

41. Vallabhaneni R, Gallagher CE, Licciardello N, Cuttriss AJ, Quinlan RF, Wurtzel ET: **Metabolite sorting of a germplasm collection reveals the hydroxylase3 locus as a new target for maize provitamin A biofortification.** *Plant Physiol* 2009, **151**:1635-1645.
42. Diretto G, Tavazza R, Welsch R, Pizzichini D, Mourgues F, Papacchioli V, Beyer P, Giuliano G: **Metabolic engineering of potato tuber carotenoids through tuber-specific silencing of lycopene epsilon cyclase.** *BMC Plant Biol* 2006, **6**:13.
43. Diretto G, Welsch R, Tavazza R, Mourgues F, Pizzichini D, Beyer P, Giuliano G: **Silencing of beta-carotene hydroxylase increases total carotenoid and beta-carotene levels in potato tubers.** *BMC Plant Biol* 2007, **7**:11.
44. Wolters AM, Uidevilligen JG, Kloosterman BA, Hutten RC, Visser RG, van Eck HJ: **Identification of alleles of carotenoid pathway genes important for zeaxanthin accumulation in potato tubers.** *Plant Mol Biol* 2010, **73**:659-671.

QTLs in the potato CHY2 and ZEP genes associated with altered carotenoid composition of tubers.

45. Romer S, Lubeck J, Kauder F, Steiger S, Adomat C, Sandmann G: **Genetic engineering of a zeaxanthin-rich potato by antisense**

- inactivation and co-suppression of carotenoid epoxidation.** *Metab Eng* 2002, **4**:263-272.
46. Verde I, Abbott AG, Scalabrin S, Jung S, Shu S, Marroni F, Zhebentyayeva T, Dettori MT, Grimwood J, Cattonaro F *et al.*: **The high-quality draft genome of peach (*Prunus persica*) identifies unique patterns of genetic diversity, domestication and genome evolution.** *Nat Genet* 2013, **45**:487-494.
The peach genome sequence.
 47. Brandi F, Bar E, Mourgues F, Horvath G, Turcsi E, Giuliano G, Liverani A, Tartarini S, Lewinsohn E, Rosati C: **Study of 'redhaven' peach and its white-fleshed mutant suggests a key role of *ccd4* carotenoid dioxygenase in carotenoid and norisoprenoid volatile metabolism.** *BMC Plant Biol* 2011, **11**:24.
A novel role for CCD4 in the control of peach flesh pigmentation.
 48. Falchi R, Vendramin E, Zanon L, Scalabrin S, Cipriani G, Verde I, Vizzotto G, Morgante M: **Three distinct mutational mechanisms acting on a single gene underpin the origin of yellow flesh in peach.** *Plant J* 2013, **76**:175-187.
Confirmation of the molecular basis for peach flesh pigmentation and demonstration of the mutational mechanisms giving rise to yellow-fleshed varieties.
 49. Yoshioka S, Aida R, Yamamizo C, Shibata M, Ohmiya A: **The carotenoid cleavage dioxygenase 4 (*cmccd4a*) gene family encodes a key regulator of petal color mutation in *Chrysanthemum*.** *Euphytica* 2012, **184**:377-387.
 50. Hai NTL, Masuda J-I, Miyajima I, Thien NQ, Mojtahedi N, Hiramatsu M, Kim J-H, Okubo H: **Involvement of carotenoid cleavage dioxygenase 4 gene in tepal color change in *Lilium brownii* var *Colchesteri*.** *J Jpn Soc Hortic Sci* 2012, **81**:366-373.
 51. Kishimoto S, Ohmiya A: **Carotenoid isomerase is key determinant of petal color of *Calendula officinalis*.** *J Biol Chem* 2012, **287**:276-285.
A molecular basis for *Calendula* petal pigmentation.
 52. Campbell R, Ducreux LJ, Morris WL, Morris JA, Suttle JC, Ramsay G, Bryan GJ, Hedley PE, Taylor MA: **The metabolic and developmental roles of carotenoid cleavage dioxygenase 4 from potato (*Solanum tuberosum* L).** *Plant Physiol* 2010, **154**:656-664.
 53. Gonzalez-Jorge S, Ha SH, Magallanes-Lundback M, Gilliland LU, Zhou A, Lipka AE, Nguyen YN, Angelovici R, Lin H, Cepela J *et al.*: **Carotenoid cleavage dioxygenase4 is a negative regulator of beta-carotene content in *Arabidopsis* seeds.** *Plant Cell* 2013, **25**:4812-4826.
A role for CCD4 in the control of β -carotene content in seeds and senescing leaves.
 54. Rodrigo MJ, Alquezar B, Alos E, Medina V, Carmona L, Bruno M, Al-Babili S, Zacarias L: **A novel carotenoid cleavage activity involved in the biosynthesis of citrus fruit-specific apocarotenoid pigments.** *J Exp Bot* 2013, **64**:4461-4478.
A novel cleavage activity for the CCD4 enzyme.
 55. Dharmapuri S, Rosati C, Pallara P, Aquilani R, Bouvier F, Camara B, Giuliano G: **Metabolic engineering of xanthophyll content in tomato fruits.** *FEBS Lett* 2002, **519**:30-34.
 56. Fujisawa M, Takita E, Harada H, Sakurai N, Suzuki H, Ohyama K, Shibata D, Misawa N: **Pathway engineering of *Brassica napus* seeds using multiple key enzyme genes involved in ketocarotenoid formation.** *J Exp Bot* 2009, **60**:1319-1332.
 57. Azadi P, Otang NV, Chin DP, Nakamura I, Fujisawa M, Harada H, Misawa N, Mii M: **Metabolic engineering of *Lilium* \times *Formolongi* using multiple genes of the carotenoid biosynthesis pathway.** *Plant Biotechnol Rep* 2010, **4**:269-280.
 58. Ha SH, Liang YS, Jung H, Ahn MJ, Suh SC, Kweon SJ, Kim DH, Kim YM, Kim JK: **Application of two bicistronic systems involving 2A and IRES sequences to the biosynthesis of carotenoids in rice endosperm.** *Plant Biotechnol J* 2010, **8**:928-938.
A comparison of different strategies for expressing multiple nuclear transgenes in tandem in a single transcript.
 59. Kim MJ, Kim JK, Kim HJ, Pak JH, Lee JH, Kim DH, Choi HK, Jung HW, Lee JD, Chung YS, Ha SH: **Genetic modification of the soybean to enhance the beta-carotene content through seed-specific expression.** *PLoS ONE* 2012, **7**:e48287.
 60. Kumar S, Hahn FM, Baidoo E, Kahlon TS, Wood DF, McMahan CM, Cornish K, Keasling JD, Daniell H, Whalen MC: **Remodeling the isoprenoid pathway in tobacco by expressing the cytoplasmic mevalonate pathway in chloroplasts.** *Metab Eng* 2012, **14**:19-28.
A transformation 'tour de force' in which a whole cytoplasmic pathway is transferred to a different organelle. Would have been of exceptional interest after a more thorough characterization of the expression and localization of the different enzymes.
 61. Nogueira M, Mora L, Enfissi EM, Bramley PM, Fraser PD: **Subchromoplast sequestration of carotenoids affects regulatory mechanisms in tomato lines expressing different carotenoid gene combinations.** *Plant Cell* 2013, **25**:4560-4579.
Novel insights on the compartmentation of the carotenoid intermediates and enzymes in metabolically engineered tomato plastids.
 62. Ytterberg AJ, Peltier JB, van Wijk KJ: **Protein profiling of plastoglobules in chloroplasts and chromoplasts. A surprising site for differential accumulation of metabolic enzymes.** *Plant Physiol* 2006, **140**:984-997.
 63. Rubio A, Rambla JL, Santaella M, Gomez MD, Orzaez D, Granell A, Gomez-Gomez L: **Cytosolic and plastoglobule-targeted carotenoid dioxygenases from *Crocus sativus* are both involved in beta-ionone release.** *J Biol Chem* 2008, **283**:24816-24825.
 64. Zhong YJ, Huang JC, Liu J, Li Y, Jiang Y, Xu ZF, Sandmann G, Chen F: **Functional characterization of various algal carotenoid ketolases reveals that ketolating zeaxanthin efficiently is essential for high production of astaxanthin in transgenic *Arabidopsis*.** *J Exp Bot* 2011, **62**:3659-3669.
 65. Huang J, Zhong Y, Sandmann G, Liu J, Chen F: **Cloning and selection of carotenoid ketolase genes for the engineering of high-yield astaxanthin in plants.** *Planta* 2012, **236**:691-699.
 66. Caroca R, Howell KA, Hasse C, Ruf S, Bock R: **Design of chimeric expression elements that confer high-level gene activity in chromoplasts.** *Plant J* 2013, **73**:368-379.
Development of a toolbox for synthetic biology in chromoplasts.
 67. Khalil AS, Collins JJ: **Synthetic biology: applications come of age.** *Nat Rev Genet* 2010, **11**:367-379.
 68. Mintz-Oron S, Meir S, Malitsky S, Ruppin E, Aharoni A, Shlomi T: **Reconstruction of *Arabidopsis* metabolic network models accounting for subcellular compartmentalization and tissue-specificity.** *Proc Natl Acad Sci U S A* 2012, **109**:339-344.
 69. Neuman H, Galpaz N, Cunningham FX Jr, Zamir D, Hirschberg J: **The tomato mutation *nxd1* reveals a gene necessary for neoxanthin biosynthesis and demonstrates that violaxanthin is a sufficient precursor for abscisic acid biosynthesis.** *Plant J* 2014, **78**:80-93.